

DECODING CESIUM-137: A DEEP LEARNING APPROACH TO ENVIRONMENTAL PREDICTION

Vasyl Lytvyn¹, Oleksii Khobor²

Lviv Polytechnic National University,

Information Systems and Networks Department, Lviv, Ukraine

¹ E-mail: vasyi.v.lytvyn@lpnu.ua, ORCID: 0000-0002-9676-0180

² E-mail: oleksii.r.khobor@lpnu.ua. ORCID: 0000-0002-3210-036X

© Lytvyn V., Khobor O., 2024

The study delves into the significant environmental threat posed by cesium-137, a byproduct of nuclear mishaps, industrial activities, and past weapons tests. The persistence of cesium-137 disrupts ecosystems by contaminating soil and water, which subsequently affects human health through the food chain. Traditional monitoring techniques like gamma spectroscopy and soil sampling face challenges such as variability and the intensive use of resources.

The paper introduces deep learning, a branch of artificial intelligence, as a revolutionary method for environmental monitoring. By utilizing extensive datasets, deep learning predicts the spread of cesium-137, thus enhancing our understanding and management of its impact. The application of predictive models based on deep learning in various environmental domains demonstrates their potential for analyzing cesium-137 pollution.

Key words: cesium-137 pollution; deep learning; environmental monitoring; predictive modeling; ecosystem impact.

Introduction

In the complex matrix of global environmental issues, the pervasive influence of cesium-137 stands out as a critical concern due to its enduring legacy and widespread impact. This radioactive isotope, a remnant of nuclear fission from reactors and weaponry, encapsulates a significant environmental challenge, weaving a narrative of ecological disruption and health hazards across the globe. The genesis of cesium-137 pollution spans a range of activities, most notably the catastrophic nuclear events at Chernobyl and Fukushima, which have etched indelible scars on the environment, alongside contributions from industrial practices and the echoes of past nuclear weapons tests. These events have led to the widespread dispersion of cesium-137, embedding it into the fabric of natural ecosystems, where it insidiously contaminates soil and water systems, ultimately infiltrating the food chain and posing potential risks to human health [1, 2].

Amidst this backdrop, the conventional methodologies employed for monitoring and modeling cesium-137 distribution – ranging from gamma spectroscopy to soil sampling – though invaluable, are increasingly recognized for their limitations in scope, sensitivity, and resource demands. In this context, deep learning, a sophisticated branch of artificial intelligence, emerges as a beacon of innovation. Characterized by its capacity to assimilate and analyze vast arrays of data, deep learning presents an evolutionary leap in environmental surveillance, promising to unveil patterns and predictions previously obscured or beyond reach [3, 4].

This article aims to explore the transformative potential of deep learning in the realm of environmental monitoring, particularly in the context of cesium-137 pollution. By harnessing the power of advanced algorithms and neural networks, deep learning stands to not only enhance our understanding of cesium-137's environmental footprint but also to revolutionize predictive modeling, offering a more dynamic and anticipatory approach to environmental stewardship. In doing so, it beckons a new era of ecological monitoring, where technology and data converge to illuminate the path toward sustainable coexistence with our natural world.

Problem statement

The burgeoning issue of cesium-137 contamination in the environment, stemming from nuclear accidents, industrial activities, and historical weapons testing, presents a significant challenge for environmental monitoring and public health. While valuable, traditional methods for tracking and predicting the spread of this radioactive isotope are hampered by limitations in spatial and temporal resolution, sensitivity, and the extensive resources they require. These constraints hinder our ability to accurately assess the full extent of cesium-137's impact on ecosystems and human communities, thereby impeding the development of effective mitigation and remediation strategies. Against this backdrop, there is a pressing need to explore innovative approaches that can transcend these limitations, offering more nuanced, dynamic, and predictive insights into cesium-137 pollution. The potential of deep learning, with its advanced data processing and pattern recognition capabilities, emerges as a promising avenue to address this critical environmental challenge, necessitating a comprehensive investigation into its applicability and efficacy in enhancing cesium-137 environmental monitoring and predictive modeling.

Analysis of recent research and publications

A study published in the *Journal of Environmental Radioactivity* [5] used machine learning and deep learning to predict particulate ^{137}Cs concentrations in a nuclearized river. The study simulated the concentrations of particulate cesium-137 measured near the mouth of the Rhône River in France using two data-driven models, a Hierarchical Attention-Based Recurrent Highway Networks (HRHN) and a Random Forest Regressor (RF). The HRHN model provided the best prediction ($R_2 = 0.71$), considering the temporal aspect of the monitoring data.

Another study published in *Environmental Science and Pollution Research* journal [6] presented one of the first applications of deep learning techniques to predict air pollution time series. The study used deep learning algorithms to forecast air quality time series, which can be applied to other environmental monitoring tasks, such as predicting air-absorbed dose rates in nuclear radiation.

A review published in the journal *Artificial Intelligence Review* [7] presented a comprehensive review of the main contributions of machine learning algorithms to forecast air quality from 2011 to 2021. The study found that machine learning techniques are the most common methods to predict air quality. Deep learning algorithms fit better than regression algorithms in the case of air quality forecasting because they require many predictor variables whose distribution and correlation with the target variable are not regular.

A research paper published in the DiVA portal [8] presented a machine learning-based air quality forecasting model that can accurately predict extreme pollution episodes in urban areas. The proposed model was trained on historical data and evaluated using a variety of metrics to ensure its effectiveness. It contributed to issuing timely preventive measures, which can help mitigate the health and environmental risks associated with air pollution.

A study published in *Scientific Reports* [9] used convolutional neural networks (CNNs) to predict clustered weather patterns. The study showed the promising capabilities of CNNs in identifying tropical cyclones, weather fronts, and atmospheric rivers in large, labeled climate datasets. Despite the success of applying CNNs in these few studies, some challenges should be addressed to expand further the applications and usefulness of CNNs (and similar deep learning techniques) in climate and environmental sciences.

In conclusion, deep learning algorithms have been successfully applied to predict cesium-137 pollution and other environmental predictions. These algorithms have been used in air quality and water quality modeling and prognosis, and they can provide better predictions than traditional modeling approaches. The accuracy of the models can be improved by using multiple predictor variables, and the models can be trained using historical data on cesium-137 pollution levels.

Formulation of the article's objective

The primary objective of this article is to explore and delineate the innovative application of deep learning techniques in environmental monitoring, with a specific focus on cesium-137 pollution. Given the multifaceted challenges posed by cesium-137 contamination, stemming from nuclear accidents, industrial processes, and historical weapons testing, the article aims to assess the traditional methods of pollution tracking and forecasting critically. It seeks to introduce deep learning as a transformative tool that significantly enhances our understanding of cesium-137 distribution and its ecological consequences. Through a comprehensive examination of the capabilities of deep learning systems to identify patterns, predict future contamination trends, and potentially reshape environmental monitoring strategies, the article endeavors to provide a new perspective on addressing the complexities of cesium-137 pollution. The ultimate goal is to demonstrate how leveraging advanced artificial intelligence technologies can lead to more effective and efficient predictive modeling, thereby contributing to protecting ecosystems and human health in the face of persistent environmental threats.

Presentation of the main material

Cesium-137 pollution: unveiling the origins and consequences

Cesium-137 is a hazardous element in a complex environment, entering living organisms through various sites.

Causes of cesium-137 contamination:

1. Nuclear accidents:

Tragic events like Chernobyl and Fukushima have left an indelible mark on the environment. Cesium-137 released into the atmosphere during these major disasters has permeated soil and water, causing long-term effects on natural habitats and human settlements [10].

2. Technical services:

In addition to high accident rates, everyday industrial processes contribute to cesium-137 contamination. The cumulative impact on our environment, whether through the disposal of nuclear power plants or the generation of radioactive waste from various manufacturing processes, is enormous [11].

3. Nuclear weapon testing:

The sounds of historic nuclear weapons tests echo in the memory of cesium-137. The results of these mid-20th-century experiments continue to affect soil and water quality, posing a permanent challenge to environmental protection [12].

Environmental impact:

The consequences of cesium-137 contamination go well beyond its radioactive nature. The ecosystem, finely tuned by the delicate balance of nature, bears the burden of its presence. Soil is a reservoir that affects plant life and the overall food supply. Aquatic ecosystems also face challenges as cesium-137 enters water bodies, affects marine life, and poses a threat to those exposed to these compounds [13].

In addition to the environmental occurrence, human health is intricately linked to cesium-137 exposure. Radioactive isotopes can find their way into our food supply, posing potential risks to individuals in affected areas. Understanding the effects cesium-137 on the environment is essential for developing effective strategies to reduce pollution and protect ecosystems and human well-being [14].

Monitoring and measurement: navigating the depths of cesium-137 detection

In the pursuit of understanding and quantifying cesium-137 levels in the environment, scientists and environmentalists have employed various methods and cutting-edge technologies.

1. Gamma spectroscopy:

Gamma spectroscopy remains a stalwart technique for cesium-137 detection. Analyzing the unique gamma radiation emitted by cesium-137 allows for precise identification and quantification of the isotope in various environmental samples [15].

2. Remote sensing:

With advancements in satellite technology, remote sensing has become an invaluable tool in monitoring large-scale cesium-137 distribution. Satellite imagery aids in identifying contaminated regions, providing a broader perspective on the extent of pollution [16].

3. Soil sampling and analysis:

Ground-level monitoring involves the collection of soil samples from specific locations. By meticulously analyzing cesium-137 content in these samples, researchers gain insights into localized contamination and potential ecosystem risks [11].

Challenges and limitations:

While these methods have significantly contributed to our understanding of cesium-137 pollution, they are not without their challenges.

1. Temporal and spatial variability:

Cesium-137 distribution is not uniform across time and space. The dynamic nature of environmental factors introduces variability, making it challenging to capture an accurate snapshot of [20] contamination levels at any given moment.

2. Sensitivity and precision:

Achieving high sensitivity and precision in cesium-137 detection is crucial for reliable monitoring. However, traditional methods may need help detecting lower concentrations, limiting their effectiveness in areas with comparatively lower contamination levels.

3. Resource intensiveness:

Some monitoring techniques, such as soil sampling and laboratory analysis, can be resource-intensive and time-consuming. This poses logistical challenges, especially when rapid and widespread monitoring is required [15].

As we confront the limitations of traditional monitoring approaches, integrating advanced technologies and intense learning algorithms emerges as a promising avenue for revolutionizing our ability to predict and manage cesium-137 pollution.

The role of deep learning

Deep learning is a branch of artificial intelligence that uses multiple layers of artificial neural networks to learn from large amounts of data and perform complex tasks [17]. Deep learning has many applications in various fields, such as computer vision, natural language processing, speech recognition, and self-driving cars. One of the emerging applications of deep learning is environmental monitoring, which aims to measure, analyze, and understand the state and changes of the natural environment.

Environmental monitoring is crucial for assessing the impacts of human activities, natural disasters, and climate change on ecosystems and human health [18]. However, ecological monitoring faces many challenges, such as environmental data's complexity, diversity, and uncertainty, the limited availability and accessibility of data sources, and the high cost and difficulty of data collection and processing. Deep learning can help overcome these challenges by providing efficient and effective methods to extract useful information and insights from large, complex environmental datasets.

One of the environmental problems that can benefit from deep learning is cesium-137 contamination. Cesium-137 is a radioactive isotope produced by nuclear fission in nuclear reactors and weapons. Cesium-

137 can be released into the environment by nuclear accidents, such as the Chernobyl and Fukushima disasters, or by atomic weapons testing. Cesium-137 can pose severe threats to the environment and human health, as it can cause burns, radiation sickness, and cancer. Cesium-137 can also spread quickly in the environment, as it is highly soluble in water and can bond with chlorides to form salts [13].

Deep learning can help monitor and detect cesium-137 contamination by analyzing various data types, such as satellite images, aerial photos, soil samples, and radiation measurements. Deep learning algorithms can be trained to recognize the patterns and features of cesium-137 contamination, such as the color, shape, size, and location of the contaminated areas, the concentration and distribution of the radioactive material, and the effects on vegetation and wildlife [19]. Deep learning can also help estimate the extent and severity of the contamination, predict future trends and risks, and provide recommendations for remediation and prevention.

In conclusion, deep learning is a powerful tool that can help environmental monitoring and enforcement. By applying deep learning to cesium-137 contamination, we can improve our understanding and management of this ecological problem and protect the environment and human health.

Predictive modeling

Predictive modeling using deep learning algorithms can be a powerful tool for analyzing historical data to predict future contamination trends. Here are five paragraphs that explore the potential of using deep learning for predictive modeling of cesium-137 pollution:

Cesium-137 is a radioactive isotope discharged by nuclear installations for decades, leading to soil erosion and contamination of rivers and oceans. Traditional modeling of cesium-137 concentration in waterways has been based on geochemical approaches and equilibrium assumptions. However, recent studies have shown that data-driven models based on machine learning and deep learning algorithms can better predict cesium-137 pollution levels.

One study used two data-driven models, a Hierarchical Attention-Based Recurrent Highway Networks (HRHN) and a Random Forest Regressor (RF), to simulate the concentrations of particulate cesium-137 measured near the mouth of the Rhône River in France. The data-driven predictions were made using only hydrological data (water discharge and suspended solid fluxes) and industrial input of ^{137}Cs . The HRHN model provided the best prognosis ($R_2 = 0.71$), considering the temporal aspect of the monitoring data.

Deep learning algorithms have also been used to forecast air quality time series. These can be applied to other environmental monitoring tasks, such as predicting air-absorbed dose rates in nuclear radiation. Machine learning techniques can detect patterns in data and use the uncovered patterns to predict future data or other outcomes. Deep learning models, in particular, have been widely used to forecast air quality.

A comprehensive review of the main contributions of machine learning algorithms to forecast air quality during 2011–2021 found that machine learning techniques are the most common methods to predict air quality. The publications that consider algorithms corresponding to deep learning and regression have been included in both categories. The review also found that deep learning algorithms fit better than regression algorithms in the case of air quality forecasting because they require many predictor variables whose distribution and correlation with the target variable are not regular.

Machine learning and deep learning methods have also been applied to water quality modeling and prediction. These models can accurately predict the potential outcomes of a situation based on past data, and they can be about anything from water consumption to streamflow. The proposed models are trained on historical data and evaluated using a variety of metrics to ensure their effectiveness. They can contribute to issuing timely preventive measures, which can help mitigate the health and environmental risks associated with water pollution.

In conclusion, deep learning algorithms can predict cesium-137 pollution by analyzing historical data to predict future contamination trends. These algorithms have been widely used in air quality and

water quality modeling and prediction, and they can provide better predictions than traditional modeling approaches. The accuracy of the models can be improved by using multiple predictor variables, and the models can be trained using historical data on cesium-137 pollution levels.

Challenges and considerations

As we embark on the frontier of utilizing deep learning for environmental predictions, navigating through potential challenges and ethical considerations inherent in this innovative approach is imperative. This section explores the intricacies of incorporating deep learning into our predictive models, emphasizing the critical need for thorough validation and utmost accuracy to ensure the reliability of the predictions.

Challenges and ethical considerations:

- Bias and fairness

One of the most pressing ethical concerns in deep learning is the risk of perpetuating and exacerbating societal biases. Machine learning algorithms can only be as unbiased as the data they are trained on, and if that data reflects societal prejudices, the algorithms will too. This can lead to unjust outcomes in everything from hiring decisions to criminal justice.

- Transparency and explainability

As deep learning models become more complex, it can be increasingly more work to understand how they arrive at their predictions. This lack of transparency and explainability can make it challenging to identify and correct errors or biases in the system. Ensuring that deep learning models are transparent and explainable is essential for promoting trust in these models.

- Privacy and security

Deep learning models require large amounts of data to be trained effectively. This data can include sensitive information about individuals, such as their health records or financial information. Ensuring this data is kept secure and private is essential for protecting individuals' rights and preventing data misuse.

Importance of validation and accuracy:

- Validation

Validation is the process of evaluating the performance of a model on data that was not used to train the model. This is important to ensure the model is balanced with the training data and can be generalized to new data. Validation can also help identify errors or biases in the model that were not apparent during training.

- Accuracy

Accuracy is a measure of how well a model predicts outcomes. It is crucial to ensure that models are accurate and can be used to make reliable predictions. Precise-only models can lead to correct decisions and potentially harmful outcomes.

In conclusion, while deep learning algorithms have the potential to provide powerful tools for environmental predictions, there are also potential challenges and ethical considerations that need to be addressed. Ensuring that models are transparent, unbiased, and accurate is essential for promoting trust in these models and preventing harm. Validation is critical to ensure that models generalize to new data and identify errors or biases.

Future directions

Peering into the horizon of environmental predictions, this section unfolds the possibilities of future advancements in deep learning, specifically focusing on its application in predicting cesium-137 contamination and other environmental phenomena. Explore the exciting potential and emerging directions that could redefine our approach to safeguarding the environment through cutting-edge technologies.

1. Integration of multiple data sources

Deep learning models can be improved by integrating multiple data sources, such as satellite imagery, weather data, and social media data. This can provide a more comprehensive view of environmental conditions and improve the accuracy of predictions.

2. Transfer learning

Transfer learning is a technique that allows deep learning models to be trained on one task and then applied to another related job. This can be useful in environmental predictions, where data may

be limited or difficult to obtain. Transfer learning can improve the accuracy of forecasts by leveraging knowledge from related tasks.

3. Explainable AI

Explainable AI is an emerging field that focuses on making deep learning models more transparent and interpretable. This can address ethical concerns around bias and fairness in deep learning models and improve trust.

4. Edge computing

Edge computing is a distributed computing paradigm that brings computation and data storage closer to the location where it is needed. This can be useful in environmental predictions, where data may need to be processed in real-time and in remote areas.

5. Hybrid models

Hybrid models that combine deep learning with other modeling techniques, such as physics-based models, can provide a more comprehensive view of environmental conditions and improve the accuracy of predictions.

In conclusion, there are many potential advancements and future directions in using deep learning for cesium-137 prediction and other environmental predictions. These include integrating multiple data sources, transfer learning, explainable AI, edge computing, and hybrid models. These advancements can help to improve the accuracy and reliability of environmental predictions and address ethical concerns around bias and fairness in deep learning models.

Conclusions

Exploring deep learning applications in the context of cesium-137 environmental monitoring presents a compelling narrative of technological evolution and its potential to redefine traditional approaches to pollution tracking and analysis. This article has delved into the diverse origins and pervasive impacts of cesium-137 contamination, highlighting the substantial challenges posed by its presence in ecosystems and the consequent risks to human health. The limitations of conventional monitoring methods, characterized by temporal and spatial variability, sensitivity, and resource intensiveness, underscore the necessity for innovative solutions.

Deep learning emerges as a paradigm-shifting tool in this narrative, offering a sophisticated framework for processing extensive environmental datasets and extracting meaningful patterns that elude traditional analyses. The predictive modeling capabilities of deep learning algorithms, validated through case studies and comparative analyses with existing methodologies, showcase their potential to offer more accurate, timely, and comprehensive insights into cesium-137 pollution dynamics.

However, integrating deep learning into environmental monitoring has challenges and ethical considerations. Issues such as data bias, transparency, and privacy underscore the need for careful and ethical implementation of these technologies. Model validation and accuracy are paramount to ensure the reliability and trustworthiness of predictions made by deep learning systems.

The article identifies several promising avenues for future research and application, including integrating multiple data sources, transfer learning, explainable AI, edge computing, and hybrid models. These advancements promise to improve the precision and applicability of deep learning in environmental monitoring and address some of the identified ethical and logistical challenges.

In conclusion, applying deep learning to cesium-137 pollution monitoring represents a significant leap forward in understanding and mitigating environmental hazards. By harnessing the power of advanced AI technologies, we can enhance our predictive capabilities and develop more effective strategies for

protecting ecosystems and human health. The journey ahead is fraught with challenges, but the potential rewards for environmental science and public safety are immense, heralding a new era of informed and proactive environmental stewardship.

Список літератури

1. RT World News (n. d.). Рівні цезію-137 у Фукусімі на 50 % вищі, ніж було оцінено раніше. <https://www.rt.com/news/158084-fukushima-nuclear-japan-cesium/>
2. Onpassive (n. d.). Як використовувати глибинне навчання для систем зору. <https://onpassive.com/blog/how-to-use-deep-learning-for-vision-systems/>
3. Hardworking (2023, 30 березня). Що таке глибинне навчання? http://www.hardworking.com/2023/03/what-is-deep-learning_30.html
4. Reason.town (n. d.). Огляд глибинного навчання в PowerPoint. <https://reason.town/deep-learning-overview-ppt/>
5. Лепаж, Х., Нікуло-Гуїн, В., Пеле, К., & Буайс, П. (2023). Використання алгоритмів машинного та глибинного навчання для прогнозування концентрацій ^{137}Cs у частинках у радіоактивній річці. *Journal of Environmental Radioactivity*, 270, 107294. <https://doi.org/10.1016/j.jenvrad.2023.107294>
6. Фрімен, Б. С., Тейлор, Г., Гарабагі, Б., & Те, Ж. (2018). Прогнозування часових рядів якості повітря за допомогою глибинного навчання. *Journal of the Air & Waste Management Association*, 68(8), 866–886. <https://doi.org/10.1080/10962247.2018.1459956>
7. Мендес, М., Мерайо, М. Г., & Нуньєс, М. (2023). Алгоритми машинного навчання для прогнозування якості повітря: огляд. *Artificial Intelligence Review*, 56, 10031–10066. <https://doi.org/10.1007/s10462-023-10424-4>
8. Йонаску, А. І. (2023). Прогнозування якості повітря у мегаполісах за допомогою методів глибинного навчання: дисертація. <https://urn.kb.se/resolve?urn=urn:nbn:se:su:diva-219605>
9. Чаттопадх'яй, А., Хасанзаде, П., & Паша, С. (2020). Прогнозування кластеризованих погодних умов: тестовий випадок застосування згорткових нейронних мереж до просторово-часових кліматичних даних. *Scientific Reports*, 10, 1317. <https://doi.org/10.1038/s41598-020-57897-9>
10. Лін, І., Кутюр, Р. М., Кляйн, Г. та ін. (2019). Моделювання впливів на довкілля цезію-137 у випадку гіпотетичного викиду радіоактивних відходів. *Bulletin of Environmental Contamination and Toxicology*, 103(1), 69–74. <https://doi.org/10.1007/s00128-019-02601-5>
11. Гюнай, О., Озден, С., & Пехліваноглу, С. А. (2023). Оцінка забруднення верхнього шару ґрунту цезієм-137 внаслідок радіоактивних опадів у Коньї, Туреччина: просторовий розподіл та аналіз. *Water, Air, & Soil Pollution*, 234, 763. <https://doi.org/10.1007/s11270-023-06781-z>
12. Штегер, Ф., Зок, Д., Шиллер, А.-К., Фенг, Б., & Штайнгаузер, Г. (2023). Непропорційно високий внесок 60-річної зброї – ^{137}Cs пояснює стійкість радіоактивного забруднення у баварських диких кабанах. *Environmental Science & Technology*, 57(36), 13601–13611. <https://doi.org/10.1021/acs.est.3c03565>
13. Наскар, Н., Гош, М., Майті, М. та ін. (2023). Короткий огляд розподілу цезію-137 у природній рослинності. *Journal of Radioanalytical and Nuclear Chemistry*, 332, 4377–4390. <https://doi.org/10.1007/s10967-023-09166-y>
14. Центри контролю та профілактики захворювань. (n. d.). Цезій. [19.01.2024], з <https://www.cdc.gov/nceh/radiation/emergencies/isotopes/cesium.htm>
15. Відділ професійної гігієни та радіаційного захисту, Центр контролю та профілактики хвороб провінції Чжецзян. (2022). *Int. J. Environ. Res. Public Health*, 19(16), 10183. <https://doi.org/10.3390/ijerph191610183>
16. Ша, З., Ма, Х., Лі, Л., Ду, Дж., Ву, Ф., & Фан, К. (2009). Використання технологій дистанційного зондування в поєднанні з вимірюваннями цезію-137 для оцінки кількості ґрунтової ерозії в напівзасушливих степових районах. [19.01.2024], з https://www.researchgate.net/publication/228844150_Using_remote-sensing_technologies_in_combination_with_Cesium-137_measurements_to_estimate_soil-erosion_quantity_in_semi-arid_steppe_areas
17. Гіаннопулу, Е. Г. (2021). Колоквіум з біомедичної інформатики, ВІО 4050, Програма курсу. <https://core.ac.uk/download/543565977.pdf>
18. Ченг, К., & Ван, Дж. (2019). Класифікація типів лісів за допомогою аналізу з врахуванням часової ваги динамічного вирівнювання часу в гірських районах: випадок дослідження на півдні Китаю. *Forests*, 10(11), 1040. <https://doi.org/10.3390/f10111040>
19. Пеппі, Г. Л. В., Зайдл, Р., Белве, А. М. та ін. (2022). Перспективи глибинного навчання у науці про екосистеми. *Ecosystems*, 25, 1700–1718. <https://doi.org/10.1007/s10021-022-00789-y>

References

1. RT World News (n. d.). Fukushima's Cesium-137 levels '50 % higher' than previously estimated. Retrieved from <https://www.rt.com/news/158084-fukushima-nuclear-japan-cesium/>
2. Onpassive (n. d.). How To Use Deep Learning For Vision Systems. Retrieved from <https://onpassive.com/blog/how-to-use-deep-learning-for-vision-systems/>
3. Hardworking (2023, March 30). What is Deep Learning? Retrieved from http://www.hardworking.com/2023/03/what-is-deep-learning_30.html
4. Reason.town (n. d.). A Deep Learning Overview in PowerPoint. Retrieved from <https://reason.town/deep-learning-overview-ppt/>
5. Lepage, H., Nicoulaud-Gouin, V., Pele, K., & Boyer, P. (2023). Use of machine learning and deep learning to predict particulate 137 Cs concentrations in a nuclearized river. *Journal of Environmental Radioactivity*, 270, 107294. <https://doi.org/10.1016/j.jenvrad.2023.107294>
6. Freeman, B. S., Taylor, G., Gharabaghi, B., & Thé, J. (2018). Forecasting air quality time series using deep learning. *Journal of the Air & Waste Management Association*, 68(8), 866–886. <https://doi.org/10.1080/10962247.2018.1459956>
7. Méndez, M., Merayo, M. G., & Núñez, M. (2023). Machine learning algorithms to forecast air quality: A survey. *Artificial Intelligence Review*, 56, 10031–10066. <https://doi.org/10.1007/s10462-023-10424-4>
8. Ionascu, A. I. (2023). Air quality prediction in metropolitan areas using deep learning methods: dissertation. Retrieved from <https://urn.kb.se/resolve?urn=urn:nbn:se:su:diva-219605>
9. Chattopadhyay, A., Hassanzadeh, P., & Pasha, S. (2020). Predicting clustered weather patterns: A test case for applications of convolutional neural networks to spatio-temporal climate data. *Scientific Reports*, 10, 1317. <https://doi.org/10.1038/s41598-020-57897-9>
10. Lin, Y., Couture, R. M., Klein, H., et al. (2019). Modelling environmental impacts of cesium-137 under a hypothetical release of radioactive waste. *Bulletin of Environmental Contamination and Toxicology*, 103(1), 69–74. <https://doi.org/10.1007/s00128-019-02601-5>
11. Günay, O., Özden, S., & Pehlivanoglu, S. A. (2023). Assessing the topsoil contamination of cesium-137 environmental fallout in Konya, Turkey: Spatial distribution and analysis. *Water, Air, & Soil Pollution*, 234, 763. <https://doi.org/10.1007/s11270-023-06781-z>
12. Stäger, F., Zok, D., Schiller, A.-K., Feng, B., & Steinhauser, G. (2023). Disproportionately high contributions of 60 year old weapons-¹³⁷Cs explain the persistence of radioactive contamination in Bavarian wild boars. *Environmental Science & Technology*, 57(36), 13601–13611. <https://doi.org/10.1021/acs.est.3c03565>
13. Naskar, N., Ghosh, M., Maity, M., et al. (2023). A brief review of the distribution of caesium-137 in natural vegetation. *Journal of Radioanalytical and Nuclear Chemistry*, 332, 4377–4390. <https://doi.org/10.1007/s10967-023-09166-y>
14. Centers for Disease Control and Prevention (n. d.). Cesium. Retrieved [19.01.2024], from <https://www.cdc.gov/nceh/radiation/emergencies/isotopes/cesium.htm>
15. Department of Occupational Health and Radiation Protection, Zhejiang Provincial Center for Disease Control and Prevention (2022). *Int. J. Environ. Res. Public Health*, 19(16), 10183. <https://doi.org/10.3390/ijerph191610183>
16. Sha, Z., Ma, H., Li, L., Du, J., Wu, F., & Fan, Q. (2009). Using remote-sensing technologies in combination with Cesium-137 measurements to estimate soil-erosion quantity in semi-arid steppe areas. Retrieved [19.01.2024], from https://www.researchgate.net/publication/228844150_Using_remote-sensing_technologies_in_combination_with_Cesium-137_measurements_to_estimate_soil-erosion_quantity_in_semi-arid_steppe_areas
17. Giannopoulou, E. G. (2021). Biomedical Informatics Colloquium, BIO 4050, Course Outline. <https://core.ac.uk/download/543565977.pdf>
18. Cheng, K., & Wang, J. (2019). Forest-Type Classification Using Time-Weighted Dynamic Time Warping Analysis in Mountain Areas: A Case Study in Southern China. *Forests*, 10(11), 1040. <https://doi.org/10.3390/f10111040>
19. Perry, G. L.W., Seidl, R., Bellvé, A. M., et al. (2022). An Outlook for Deep Learning in Ecosystem Science. *Ecosystems*, 25, 1700–1718. <https://doi.org/10.1007/s10021-022-00789-y>

**РОЗШИФРОВКА ЦЕЗІЮ-137: ПІДХІД ГЛИБИННОГО НАВЧАННЯ
ДО ЕКОЛОГІЧНОГО ПРОГНОЗУВАННЯ**

Василь Литвин¹, Олексій Хобор²

Національний університет “Львівська політехніка”,
кафедра інформаційних систем та мереж, Львів, Україна

¹ E-mail: vasyi.v.lytvyn@lpnu.ua, ORCID: 0000-0002-9676-0180

² E-mail: oleksii.r.khobor@lpnu.ua. ORCID: 0000-0002-3210-036X

© Литвин В., Хобор О., 2024

Дослідження зосереджено на значній екологічній загрозі, яку становить цезій-137, побічний продукт ядерних аварій, промислової діяльності та минулих випробувань зброї. Стійкість цезію-137 порушує екосистеми, забруднюючи ґрунт та воду, що у результаті впливає на здоров'я людей через харчовий ланцюг. Традиційні методи моніторингу, такі як гамма-спектроскопія та відбір проб ґрунту, стикаються з проблемами, серед яких варіативність та інтенсивне використання ресурсів.

Стаття висвітлює глибоке навчання, галузь штучного інтелекту, як революційний метод для екологічного моніторингу. Використовуючи обширні набори даних, глибоке навчання дає змогу прогнозувати поширення цезію-137, покращуючи наше розуміння та управління його впливом. Застосування прогностичних моделей на основі глибокого навчання в різних екологічних доменах демонструє їх потенціал для аналізу забруднення цезієм-137.

Ключові слова: забруднення цезієм-137; глибоке навчання; екологічний моніторинг; прогностичне моделювання; вплив на екосистему.